

On asymptotically harmonic manifolds of negative curvature

Philippe CASTILLON and Andrea SAMBUSETTI

Abstract: We study asymptotically harmonic manifolds of negative curvature, without any cocompactness or homogeneity assumption. We show that asymptotic harmonicity provides a lot of information on the asymptotic geometry of these spaces: in particular, we determine the volume entropy, the spectrum and the relative densities of visual and harmonic measures on the ideal boundary. Then, we prove an asymptotic analogue of the classical mean value property of harmonic manifolds, and we characterize asymptotically harmonic manifolds, among Cartan-Hadamard spaces of strictly negative curvature, by the existence of an asymptotic equivalent $\tau(u)e^{Er}$ for the volume-density of geodesic spheres (with τ constant in case DR_M is bounded). Finally, we show the existence of a Margulis function, and explicitly compute it, for all asymptotically harmonic manifolds.

Mathematics Subject Classifications (2010): primary 53C20.

Key words: Asymptotically harmonic manifolds, spectrum, asymptotic geometry.

Introduction

Harmonic manifolds are those Riemannian manifolds whose geodesic spheres have constant mean curvature; equivalently, such that the volume density function, in normal coordinates at any point x , only depends on the distance $d(x, \cdot)$. Another equivalent condition is that the mean-value property

$$F(x_0) = \frac{1}{\text{vol}(S_{x_0}(R))} \int_{S_{x_0}(R)} F(x) dv_{S_{x_0}(R)}$$

holds for all harmonic functions F on M (cf. [3]).

In 1944, A. Lichnerowicz conjectured (and proved in dimension 4) that the rank one symmetric spaces (denoted ROSS, in the sequel) are the only harmonic manifolds. If this was proved to be true for compact simply connected manifolds (cf. [24]) and for negatively curved Cartan-Hadamard manifolds admitting compact quotients (cf. [4] section 9.C), E. Damek and F. Ricci constructed harmonic homogeneous manifolds which are not ROSS (cf. [7]). Since then, J. Heber proved that Damek-Ricci spaces and ROSS are the only homogeneous harmonic manifolds (cf. [10]), and then further relations between harmonicity, volume growth and Gromov hyperbolicity have been studied (cf. [20, 16]).

In several of these works, an asymptotic version of harmonicity naturally appears (cf. [9, 10]) : a Cartan-Hadamard manifold M is *asymptotically harmonic* if its horospheres have constant mean curvature h . This notion was introduced by F. Ledrappier in [18], and was mainly studied in the *cocompact* case (i.e. when the space admits compact quotients). F. Ledrappier proved that, within these spaces, asymptotic harmonicity is equivalent to the condition

$\inf \sigma(\Delta) = \frac{E^2}{4}$ (where $\sigma(\Delta)$ denotes the spectrum of the Laplacian of M , and E its volume-entropy); moreover, he showed that if M is asymptotically harmonic, then $E = nh$ and $\inf \sigma(\Delta) = \frac{n^2 h^2}{4}$. It was then proved (as a consequence of the work of Y. Benoist, P. Foulon and F. Labourie on the geodesic flow of asymptotically harmonic spaces [2, 9] and the characterization of locally symmetric spaces by their volume entropy due to G. Besson, G. Courtois and S. Gallot [4]) that the ROSS are the only asymptotically harmonic manifolds among cocompact, negatively curved Cartan-Hadamard spaces. On the other hand, in [6] necessary and sufficient conditions are given in order that a *homogeneous*, negatively curved Cartan-Hadamard manifold is asymptotically harmonic; however, as far as the authors know, the problem whether any asymptotically harmonic manifold is a ROSS or a Damek-Ricci space is still open in this class. Recently, it was also proved that, in dimension 3, the only asymptotically harmonic Cartan-Hadamard manifold of strictly negative curvature is the hyperbolic space (cf. [11, 23]).

The aim of this paper is to show that, for Cartan-Hadamard manifolds of strictly negative curvature of any dimension, even without any cocompactness or homogeneity assumption, asymptotic harmonicity provides a lot of information on the asymptotic geometry. In view of [18], we are naturally interested in the volume entropy, the spectrum and the relations between visual and harmonic measures on the ideal boundary of a general asymptotically harmonic manifold. In particular, in section §3, we show rigidity of Cartan-Hadamard asymptotically harmonic manifolds under suitable curvature bounds (*Corollary 3.7*), we determine the volume entropy and the spectrum (cf. *Theorems 3.3 & 3.4*) and, when the curvature is negatively pinched, we find sharp upper and lower bounds for the volume-growth of the horospheres (*Theorem 3.8* and ff. *Remarks 3.9 & 3.10*). Moreover, we prove an asymptotic analogue of the classical mean-value property holding on harmonic manifolds (*Theorem 3.11*). In section §4, we characterize asymptotically harmonic manifolds as those manifolds whose volume form, in normal coordinates, is asymptotically equivalent to a function $\tau(u)e^{ER}$, for some positive function τ on SM (*Theorem 4.1*); then, we show that the function τ is constant if DR_M (the derivative of the Riemann tensor) is bounded (*Proposition 4.3(ii)*). In §5 we prove the existence of a Margulis function (*Proposition 5.2*), we explicitly compute it for all asymptotically harmonic manifolds, and we find the relative densities of visual and harmonic measures on the ideal boundary (*Proposition 5.1*); we also show that they coincide when DR_M is bounded. This result is to compare to what is known in the cocompact and homogeneous cases, where coincidence of two of the three natural families of measures on the ideal boundary (visual, harmonic and Patterson-Sullivan measures) forces, respectively in the two cases, symmetry and asymptotic harmonicity of the manifold (cf. [18, 19, 27, 26, 6]); unfortunately, a similar characterization for general asymptotically harmonic Cartan-Hadamard manifolds is still missing.

The main tools we use are a comparison lemma for the second fundamental forms of two tangent spheres, which is proved in section 2, and the Riccati equation. The first section is devoted to notations and preliminary results.

We thank professor S. Gallot for his suggestions and encouragement, and professor G. Knieper for explaining us the expression of the function τ in terms of Jacobi tensors.

1 Notations

Unless otherwise stated, throughout all the paper (M, g) will always be a *Cartan-Hadamard manifold* (CH-manifold, for short) of dimension $n+1$, i.e. a complete, simply connected Riemannian manifold with nonpositive curvature.

The *ideal boundary* of M , denoted $\partial_\infty M$, is the set of equivalent classes of geodesic rays, γ and σ being equivalent if $\sup\{d(\gamma(t), \sigma(t)) \mid t \geq 0\} < \infty$ (cf. [5] definition II.8.1). For $\xi \in \partial_\infty M$, $\lim_{t \rightarrow +\infty} \gamma(t) = \xi$ will mean that ξ is the equivalence class defined by γ . The cone topology turn $M \cup \partial_\infty M$ into a compact manifold with boundary (cf [5] definition II.8.6).

For $\xi \in \partial_\infty M$ and $x \in M$, the *Busemann function* $b_{\xi, x}$, centered at ξ and vanishing at x , is defined by $b_{\xi, x}(y) = \lim_{t \rightarrow +\infty} (d(y, \gamma(t)) - t)$, where γ is the unique geodesic such that $\gamma(0) = x$ and $\lim_{t \rightarrow +\infty} \gamma(t) = \xi$. Two Busemann functions centered in the same point at infinity differ from a constant ; in many situations, we only need to know the Busemann functions up to a constant, and we shall note b_ξ some Busemann function centered in ξ . Busemann functions are Lipschitz and, on CH-manifolds, they are at least C^2 , cf. [12].

The *horospheres* centered in $\xi \in \partial_\infty M$ are the level hypersurfaces of b_ξ : $H_\xi(t) = \{x \in M \mid b_\xi(x) = t\}$; we shall also use the convenient notation $H_\xi(x)$ for the horosphere centred at ξ and passing through x . As the Busemann functions are limit of distance functions, the horospheres centered in ξ are (locally) limit of spheres whose centers tends to ξ . Since $|\nabla b_\xi| = 1$ and the gradient lines of b_ξ are the geodesics γ such that $\lim_{t \rightarrow -\infty} \gamma(t) = \xi$, we can define the *inner* unit vector field of horospheres centred at ξ as $\nu = -\nabla b_\xi$ (i.e. ν points towards the center ξ of the horosphere).

For a general hypersurface N of M , $\vec{\mathcal{A}}^N$ denotes its *second* (vector valued) *fundamental form* ; that is, for $u, v \in T_x N$, $\vec{\mathcal{A}}^N(u, v)$ is the component of $D_u^M V$ normal to N , where D^M is the connection of M and V extends v in a neighborhood of x . Associated to the choice of a unit normal vector field ν to N we then have the *second* (scalar) *fundamental form* $\mathcal{A}^N = \langle \vec{\mathcal{A}}^N, \nu \rangle$ and the *shape operator* $A^N \in \text{End}(T_x N)$, defined by $\langle A^N u, v \rangle = \langle D_u^M \nu, v \rangle = -\langle \vec{\mathcal{A}}^N(u, v), \nu \rangle$. The *mean curvature vector* of N at x is $\vec{h}^N(x) = \frac{1}{n} \text{Tr} \vec{\mathcal{A}}^N(x)$, while the (scalar) mean curvature, associated with ν , is $h^N = \langle \vec{h}^N, \nu \rangle$.

A manifold M is called *asymptotically harmonic* if all its horospheres have constant mean curvature h . The curvature of M being nonpositive, the horospheres are convex and we have $f \geq 0$ when choosing ν pointing to the center of the horosphere.

Hessian and Laplacian of Busemann functions

The second fundamental form naturally appears when restricting a function to a submanifold :

Proposition 1.1. *Let $i : N \rightarrow M$ be an isometric immersion, let $F : M \rightarrow \mathbb{R}$ be a smooth function and let $f = F|_N$ be its restriction to N .*

For all $x \in N$ and all $u, v \in T_x N$ we have

$$(\text{Hess}^N f)(u, v) = (\text{Hess}^M F)(u, v) + \langle \nabla^M F, \vec{\mathcal{A}}^N(u, v) \rangle$$

Proof. The proof is standard. □

As a consequence, the Hessian of the Busemann function is given by the second scalar fundamental form of its horospheres, with respect to the inner normal vector field ν ; taking the trace we get $\Delta b_\xi(y) = -\text{Tr}(\text{Hess}_y b_\xi) = -nh_\xi(y)$, where $h_\xi(y)$ is the mean curvature at y of the horosphere centered in ξ passing through y , with respect to ν . (Similarly, the second fundamental form of spheres is the Hessian of the distance function to the center and the Laplacian of the distance from a point x gives the mean curvature of the spheres centered in x).

It follows, by the regularity theory of solutions of elliptic equations, that for asymptotically harmonic manifolds Busemann functions and horospheres are at least as regular as the metric (whereas they are known to be real analytic on harmonic manifolds, cf. [21]). Moreover, it is then straightforward to check that, for any asymptotically harmonic manifold M with horospheres of mean curvature h , the function $f(y) = e^{-nh_\xi(y)}$ is harmonic.

The Riccati equation

Let $\xi \in \partial_\infty M$ and γ be a geodesic such that $\lim_{t \rightarrow -\infty} \gamma(t) = \xi$. For each t , let $A_\xi(t)$ be the shape operator of the horosphere centered in ξ passing through $\gamma(t)$, with respect to the inner unit vector field $\nu = -\nabla b_\xi = -\gamma'(t)$; this family of operators satisfies the Riccati equation (cf. [13] §1.3):

$$A'_\xi(t) + A_\xi^2(t) + R_M(\dot{\gamma}(t), \cdot)\dot{\gamma}(t) = 0 \quad (1.1)$$

where R_M is the Riemann tensor of M .

2 Comparison of spheres on CH-manifolds

In the sequel, we note $\mathbb{M}^n(-a^2)$ the simply connected Riemannian manifold with constant sectional curvature $-a^2$, and we shall note C_a and $\cot_a$ the functions defined by:

$$C_a(s) = \begin{cases} \frac{1}{a^2}(\cosh(as) - 1) & \text{if } a > 0 \\ \frac{s^2}{2} & \text{if } a = 0 \end{cases} \quad \text{and} \quad \cot_a(s) = \begin{cases} a \coth(as) & \text{if } a > 0 \\ \frac{1}{s} & \text{if } a = 0 \end{cases}$$

Comparison of triangles

When assuming a sectional curvature upper bound $K_M \leq -a^2$ for M , the classical Toponogov theorem (cf. [13]) implies that, given two edges of a triangle in M with angle α at the common vertex, then the third edge is larger than the one of a triangle in the model space $\mathbb{M}^2(-a^2)$ with the same lengths for the first two edges and the same angle at the common vertex. The following lemma is a slight modification of this result, where we compare the ratio of (some function of) the lengths of the third edge and of an “intermediate edge”.

Lemma 2.1 (Triangle comparison with curvature upper bound). *Let M be a CH-manifold with $K_M \leq -a^2 \leq 0$. Let (xyz) and $(\tilde{x}\tilde{y}\tilde{z})$ be triangles in M and $\mathbb{M}^2(-a^2)$ respectively, such that $r_1 = d(x, y) = d(\tilde{x}, \tilde{y})$, $r_2 = d(x, z) = d(\tilde{x}, \tilde{z})$ and $\alpha = \angle_x(y, z) = \angle_{\tilde{x}}(\tilde{y}, \tilde{z})$. Moreover, for $\theta \in]0, 1[$ let p, q and $\tilde{p}, \tilde{q}$ be respectively the points on the geodesic segments xy, xz and $\tilde{x}\tilde{y}, \tilde{x}\tilde{z}$ such that $d(x, p) = d(\tilde{x}, \tilde{p}) = \theta r_1$ and $d(x, q) = d(\tilde{x}, \tilde{q}) = \theta r_2$ (cf. figure 1). Then:*

$$\frac{C_a(d(y, z))}{C_a(d(p, q))} \geq \frac{C_a(d(\tilde{y}, \tilde{z}))}{C_a(d(\tilde{p}, \tilde{q}))} = F_a(r_1, r_2, \alpha, \theta).$$

Remark 2.2. By the cosine formula in $\mathbb{M}^2(-a^2)$ (cf. [5] proposition I.2.7) we know that the right-hand side of the above inequality only depends on the lengths r_1, r_2, α and θ , whence the existence of the function F_a .

When $a = 0$ we have $F_0 = \frac{1}{\theta^2}$ and lemma 2.1 is a direct consequence of the convexity of the distance function in $CAT(0)$ -spaces (cf. [5] proposition II.2.2).

When $a > 0$ we find :

$$F_a(r_1, r_2, \alpha, \theta) = \frac{\cosh(ar_1) \cosh(ar_2) - \sinh(ar_1) \sinh(ar_2) \cos(\alpha) - 1}{\cosh(a\theta r_1) \cosh(a\theta r_2) - \sinh(a\theta r_1) \sinh(a\theta r_2) \cos(\alpha) - 1}.$$

An important point in the proof of lemma 2.1 is that, whenever $\theta \leq 1$, the function F_a is nondecreasing with respect to α .

Proof of lemma 2.1. First consider a comparison triangle $(\bar{x}\bar{y}\bar{z})$ in $\mathbb{M}^2(-a^2)$, that is such that $d(\bar{x}, \bar{y}) = r_1$, $d(\bar{x}, \bar{z}) = r_2$, and $d(\bar{y}, \bar{z}) = d(y, z)$. Define $\bar{p}, \bar{q}$ to be the points on the geodesic segments $\bar{x}\bar{y}$ and $\bar{x}\bar{z}$ respectively, such that $d(\bar{x}, \bar{p}) = \theta r_1$, $d(\bar{x}, \bar{q}) = \theta r_2$, and let $\bar{\alpha} = \angle_{\bar{x}}(\bar{y}, \bar{z})$. By Toponogov theorem, we have $d(\bar{p}, \bar{q}) \geq d(p, q)$ and $\bar{\alpha} \geq \alpha$. Using these inequalities and remark 2.2 we




Figure 1: comparison triangles

have

$$\frac{C_a(d(y, z))}{C_a(d(p, q))} \geq \frac{C_a(d(y, z))}{C_a(d(\bar{p}, \bar{q}))} = F_a(r_1, r_2, \bar{\alpha}, \theta) \geq F_a(r_1, r_2, \alpha, \theta) = \frac{C_a(d(\tilde{y}, \tilde{z}))}{C_a(d(\tilde{p}, \tilde{q}))}$$

□

A similar inequality holds for CH-manifolds with curvature lower bound:

Lemma 2.3 (Triangle comparison with curvature lower bound). *Let M be a CH-manifold with $K_M \geq -b^2$. Let (xyz) and $(\tilde{x}\tilde{y}\tilde{z})$ be triangles in M and $\mathbb{M}^2(-b^2)$ respectively, such that $r_1 = d(x, y) = d(\tilde{x}, \tilde{y})$, $r_2 = d(x, z) = d(\tilde{x}, \tilde{z})$ and $\alpha = \angle_x(y, z) = \angle_{\tilde{x}}(\tilde{y}, \tilde{z})$. Moreover, for $\theta \in]0, 1[$ let p, q and $\tilde{p}, \tilde{q}$ be*

respectively the points on the geodesic segments xy , xz and $\tilde{x}\tilde{y}$, $\tilde{x}\tilde{z}$ such that $d(x, p) = d(\tilde{x}, \tilde{p}) = \theta r_1$ and $d(x, q) = d(\tilde{x}, \tilde{q}) = \theta r_2$. Then:

$$\frac{C_b(d(y, z))}{C_b(d(p, q))} \leq \frac{C_b(d(\tilde{y}, \tilde{z}))}{C_b(d(\tilde{p}, \tilde{q}))} = F_b(r_1, r_2, \alpha, \theta).$$

Proof. The proof is similar to that of lemma 2.1. Toponogov theorem gives $d(\bar{p}, \bar{q}) \leq d(p, q)$ and $\bar{\alpha} \leq \alpha$, and by the monotonicity of the function F_b we get

$$\frac{C_b(d(y, z))}{C_b(d(p, q))} \leq \frac{C_b(d(y, z))}{C_b(d(\bar{p}, \bar{q}))} = F_b(r_1, r_2, \bar{\alpha}, \theta) \leq F_b(r_1, r_2, \alpha, \theta) = \frac{C_b(d(\tilde{y}, \tilde{z}))}{C_b(d(\tilde{p}, \tilde{q}))}$$

□

Comparison of spheres

Let $S_x(r)$ and $S_y(R)$ be two geodesic spheres in M , with $r < R$, tangent at some point z , with $S_x(r)$ internal to $S_y(R)$. Let $\vec{\mathcal{A}}_x$ and $\vec{\mathcal{A}}_y$ (resp. $\mathcal{A}_x, \mathcal{A}_y$) be the second, vector-valued (resp. scalar) fundamental forms of $S_x(r)$ and $S_y(R)$, and let ν be the common inner unit normal vector at z . We will now compare the two second fundamental forms $\mathcal{A}_x$ and $\mathcal{A}_y$.

Let $u \in T_z S_x(r)$ be a unitary vector, and let $c_u(s)$ be the geodesic of $S_x(r)$ with initial tangent vector u . Denote by r_x and r_y the distance functions to x and y respectively, and let $r_y(s) = r_y(c_u(s))$ be the restriction of the function r_y to the curve c_u . Applying proposition 1.1 to c and r_y we find

$$r_y''(0) = (\text{Hess}^M r_y)(u, u) + \langle \nabla r_y, \vec{\mathcal{A}}_x(u, u) \rangle,$$

and, since $\text{Hess}^M r_y$ gives the second fundamental form of $S_y(R)$ w.r. to ν ,

$$r_y''(0) = \mathcal{A}_y(u, u) - \mathcal{A}_x(u, u) \quad (2.1)$$

But $r_y''(0) \leq 0$ as z is the maximum of r_y on $S_x(r)$, thus at the point z we have $\mathcal{A}_y \leq \mathcal{A}_x$ which means that $S_x(r)$ is “more curved” than $S_y(R)$. Using the above comparison lemmas for triangles, we get sharper comparison estimates for the tangent spheres :

Lemma 2.4. *Let (M, g) be a CH-manifold with $K_M \leq -a^2$. With the above notations, the second fundamental forms of $S_x(r), S_y(R)$ at the tangent point z satisfy:*

$$0 \leq \mathcal{A}_x - \mathcal{A}_y \leq (\cot_a r - \cot_a R) g$$

Moreover, if we assume $-b^2 \leq K_M$ then at the tangent point z we also have:

$$(\cot_b r - \cot_b R) g \leq \mathcal{A}_x - \mathcal{A}_y$$

Remark 2.5. These estimates are optimal, since they are equalities when M has, respectively, constant curvature $-a^2$ or $-b^2$.

Proof. We only consider the case $a > 0$; when $a = 0$, the proof is similar (just replace the hyperbolic laws by the Euclidean ones) and is left to the reader.

As before, let $u \in T_z S_x(r)$ be a unitary vector, let $c(s)$ be the geodesic of $S_x(r)$ with initial tangent vector u and let $r_y(s)$ be the restriction of the function r_y to the curve c . For $s > 0$ we consider (cf. figure 2) :

- the angle $\alpha(s)$ between ∇r_y and ∇r_x at $c(s)$;
- the angle $\beta(s)$ between the geodesic lines from y to z and from y to $c(s)$;
- $\theta = \frac{R-r}{R}$ and the point $x(s)$ of the geodesic from y to $c(s)$ such that $d(y, x(s)) = \theta d(y, c(s))$;

so $r'_y(s) = \langle \nabla r_y, \dot{c}(s) \rangle = -\sin(\alpha(s))$. Using Toponogov theorem for the triangle




Figure 2: Comparing the second fundamental forms of tangent spheres

$(c(s)x(s)x)$ and the law of cosine in $\mathbb{M}^2(-a^2)$ we get

$$\begin{aligned} \cosh(ad(x, x(s))) &\geq \cosh(ar) \cosh(a(1-\theta)r_y(s)) \\ &\quad - \sinh(ar) \sinh(a(1-\theta)r_y(s)) \cos(\alpha(s)) \\ &\geq 1 + \sinh(ar) \sinh(a(1-\theta)r_y(s))(1 - \cos(\alpha(s))) \end{aligned} \quad (2.2)$$

On the other hand, lemma 2.1 applied to the triangle $(yzc(s))$ implies that

$$\cosh(ad(x, x(s))) - 1 \leq \frac{\cosh(ad(z, c(s))) - 1}{F_a(R, r_y(s), \beta(s), \theta)} \quad (2.3)$$

which, plugged in (2.2), yields :

$$1 - \cos(\alpha(s)) \leq \frac{\cosh(ad(z, c(s))) - 1}{\sinh(ar) \sinh(a(1 - \theta)r_y(s)) F_a(R, r_y(s), \beta(s), \theta)} \quad (2.4)$$

We divide by s^2 and pass to the limit for $s \rightarrow 0$ in (2.4) : as $r'_y(s)^2 = \sin^2 \alpha(s)$ and $r'_y(0) = 0$, we have $\lim_{s \rightarrow 0} \frac{1 - \cos \alpha(s)}{s^2} = \frac{1}{2} \lim_{s \rightarrow 0} \left(\frac{r'_y(s)}{s} \right)^2 = \frac{1}{2} r''_y(0)^2$; then, notice that $\frac{d(z, c_u(s))}{s} \rightarrow 1$ and that, as $r_y(s) - R = O(s^2)$ and $\beta(s) = O(s)$, we have $\lim_{s \rightarrow 0} F_a(R, r_y(s), \beta(s), \theta) = \frac{\sinh^2(aR)}{\sinh^2(a(R-r))}$. So from (2.4) we get

$$|r''_y(0)| \leq \frac{a \sinh(a(R-r))}{\sinh(ar) \sinh(aR)} = a(\coth(ar) - \coth(aR))$$

By (2.1), as $r''_y(0) \leq 0$ we deduce $\mathcal{A}_x(u, u) - \mathcal{A}_y(u, u) \leq \cot_a r - \cot_a R$.

Consider now the curvature lower bound $-b^2 \leq K_M$. By Toponogov theorem and the law of cosine, equation (2.2) becomes

$$1 - \cos(\alpha(s)) \geq \frac{\cosh(bd(x, x(s))) - 1}{\sinh(br) \sinh(b(1 - \theta)r_y(s))} + \frac{1 - \cosh(br(1 - \frac{r_y(s)}{R}))}{\sinh(br) \sinh(b(1 - \theta)r_y(s))} \quad (2.5)$$

while lemma 2.3 implies

$$\cosh(bd(x, x(s))) - 1 \geq \frac{\cosh(bd(z, c(s))) - 1}{F_b(R, r_y(s), \beta(s), \theta)}, \quad (2.6)$$

which plugged in (2.5) yields

$$1 - \cos(\alpha(s)) \geq \frac{\cosh(bd(z, c(s))) - 1}{\sinh(br) \sinh(b(1 - \theta)r_y(s)) F_b(R, r_y(s), \beta(s), \theta)} + \frac{1 - \cosh(br(1 - \frac{r_y(s)}{R}))}{\sinh(br) \sinh(b(1 - \theta)r_y(s))} \quad (2.7)$$

Dividing by s^2 and letting $s \rightarrow 0$ as before, we get $(\cot_b r - \cot_b R)g \leq \mathcal{A}_x - \mathcal{A}_y$. $\square$

In the sequel, we will be mainly interested in the second fundamental form of horospheres. We will use a result similar to lemma 2.4, where the sphere $S_y(R)$ is replaced by a horosphere:

Lemma 2.6. *Let (M, g) be a CH-manifold with $K_M \leq -a^2$.*

Let $S_x(r)$ and $H_\xi(z)$ be respectively a sphere and a horosphere tangent at a point z , with $S_x(r)$ internal to the horosphere. Let $\mathcal{A}_x, \mathcal{A}_\xi$ be the second fundamental forms of $S_x(r), H_\xi(z)$ with respect to be the common inner unit normal vector at z . Then, at the tangent point z we have :

$$0 \leq \mathcal{A}_x - \mathcal{A}_\xi \leq (\cot_a r - a)g \quad (2.8)$$

Moreover, if we assume $-b^2 \leq K_M$, then at the tangent point z we also have:

$$(\cot_b r - b)g \leq \mathcal{A}_x - \mathcal{A}_\xi$$

This result can be obtained in two different ways: taking limits, in the inequalities of lemma 2.4, for y tending to ξ along the geodesic $\exp_z(t\nu)$, or following the same proof with the Busemann function b_ξ in place of r_y . The proof is left to the reader.

3 Asymptotically harmonic CH-manifolds

In this section, M will always be an asymptotically harmonic CH-manifold with horospheres of constant mean curvature h .

3.1 The entropy and the spectrum

We are interested here in two invariants of the manifold M : the volume entropy and the spectrum. The entropy is determined by the behaviour of the volume of balls whose second derivative (with respect to the radius) is given, in turns, by the mean curvature of the spheres. On the other hand, the spectrum can be determined by using special functions whose Laplacian has a nice behaviour; in our case, the distance function, whose Laplacian is again given by the mean curvature of spheres (see discussion in §1).

For points x and y in M , let $\vec{h}_x(y)$ be the mean curvature vector at y of the sphere $S_x(d(x, y))$, and $h_x(y) = -\langle \vec{h}_x(y), \nabla r_x \rangle$. Notice that, as $K_M \leq 0$, balls and horoballs are convex, so both h and $h_x(y)$ are non-negative.

Lemma 3.1. *Let M^{n+1} be an asymptotically harmonic CH-manifold. For all $x \in M$ and $r > 0$, the sphere $S_x(r)$ satisfies*

$$\forall y \in S_x(r) \quad h \leq h_x(y) \leq h + \frac{1}{r}.$$

Proof. From lemma 2.6 we have

$$\mathcal{A}_\xi(u, u) \leq \mathcal{A}_x(u, u) \leq \mathcal{A}_\xi(u, u) + \frac{1}{r}|u|^2$$

where $\mathcal{A}_x$ and $\mathcal{A}_\xi$ are, respectively, the second fundamental forms of $S_x(r)$ and of the horosphere $H_\xi(y)$, tangent to $S_x(r)$ at y . Taking the trace on an orthonormal basis gives the result. $\square$

We fix $x \in M$. For $r > 0$, let $B_x(r)$ be the ball of radius r centered in x , and $V(r) = \text{Vol}(B_x(r))$ the growth function. The *entropy* of M is defined by

$$E = \limsup_{r \rightarrow \infty} \frac{1}{r} \log V(r).$$

A first consequence of asymptotic harmonicity is the following linear isoperimetric inequality :

Proposition 3.2. *Let M^{n+1} be an asymptotically harmonic CH-manifold. For any domain $\Omega \subset M$ with smooth boundary $\partial\Omega$ we have $nh \text{Vol}(\Omega) \leq \text{vol}(\partial\Omega)$.*

Proof. Fix some $\xi \in \partial_\infty M$. Since $-\Delta b_\xi = nh$, integrating by parts on Ω the function $-\Delta b_\xi$ gives the result. $\square$

Theorem 3.3. *Let M^{n+1} be an asymptotically harmonic CH-manifold. The entropy of M is $E = nh$.*

Proof. By the co-area formula we have $V'(r) = \text{vol}(S_x(r))$, and by proposition 3.2 we get $nhV(r) \leq V'(r)$. Integrating this inequality we get $V(r) \geq Ae^{nhr}$ for some constant A , so that the entropy is bounded below by nh .

Now, the second derivative of V is given by $V''(r) = n \int_{S_x(r)} h_x(y) dv_r(y)$ where dv_r is the volume form of $S_x(r)$. Choose $\varepsilon > 0$ and let $r_0 = \frac{1}{\varepsilon}$. By lemma 3.1, we have $V''(r) \leq n(h + \varepsilon)V'(r)$ for any $r \geq r_0$. Integrating this inequality between r_0 and r , yields $V'(r) \leq Ae^{n(h+\varepsilon)r}$ for some constant A . Integrating once again between r_0 and r , we get $V(r) \leq B + Ce^{n(h+\varepsilon)r}$, which implies that $E \leq n(h + \varepsilon)$. Since ε is arbitrarily small, this concludes the proof. $\square$

Theorem 3.4. *Let M^{n+1} be an asymptotically harmonic CH-manifold.*

The spectrum of the Laplacian of M is $\sigma(\Delta) = [\frac{n^2 h^2}{4}, +\infty)$

Proof. By proposition 3.2 and Cheeger's inequality, we have $\sigma(\Delta) \subset [\frac{n^2 h^2}{4}, +\infty)$.

Conversely, we choose $x \in M$ and consider the distance function r_x to x . Since the Laplacian of r_x is given by the mean curvature of spheres, we have

$$\sup_{y \in M \setminus B_x(R)} \{ |\Delta r_x(y) - nh| \} \leq \frac{n}{R} \quad (3.1)$$

Using (3.1) and the fact that $|\nabla r_x| = 1$, we can follow the method initiated by H. Donnelly to determine the essential spectrum (cf. [8]) : for each $\lambda > \frac{n^2 h^2}{4}$ we use radial functions to construct sequences satisfying Weyl's criterion for λ (cf. [22] theorem VII.12 p. 237). See for example [17] theorem 1.2 for a general result, whose hypotheses are satisfied by the function r_x . $\square$

Remark 3.5. From theorems 3.3 and 3.4 we deduce $\inf\{\sigma(\Delta)\} = \frac{E^2}{4}$.

For cocompact negatively curved manifolds, this equality is equivalent to the asymptotic harmonicity (cf. [18] theorem 1). But, in the general case, it is easy to construct manifolds satisfying this inequality, which are not asymptotically harmonic. For example, the conclusions of theorems 3.3 and 3.4 hold true for any Cartan-Hadamard manifold with curvature less than $-h^2$ and tending to $-h^2$ at infinity.

3.2 Rigidity

Consider the second fundamental form $\mathcal{A}_\xi$ of a horosphere H_ξ , and let $\lambda_1, \dots, \lambda_n$ be the principal curvatures of H_ξ at some point x , with respect to the inner unit normal of H_ξ . If M satisfies the curvature upper bound $K_M \leq -a^2$, then it is well known that $\lambda_i \geq a$ (cf [13]). Therefore we get

$$n^2 h^2 = \left(\sum_i \lambda_i \right)^2 = \sum_i \lambda_i^2 + 2 \sum_{i < j} \lambda_i \lambda_j \geq |\mathcal{A}_\xi|^2 + n(n-1)a^2,$$

and

$$|\mathcal{A}_\xi|^2 \leq n^2 h^2 - n(n-1)a^2. \quad (3.2)$$

When assuming a curvature lower bound $K_M \geq -b^2$, a similar argument gives

$$|\mathcal{A}_\xi|^2 \geq n^2 h^2 - n(n-1)b^2. \quad (3.3)$$

Now, as the mean curvature is the same for all horospheres, taking the trace of Riccati equation (1.1) gives $|\mathcal{A}_\xi|^2 + \text{Ric}_M(u, u) = 0$ for any $u \in SM$, for the second fundamental form $\mathcal{A}_\xi$ of a horosphere tangent to $u^\perp$. Therefore we get :

Proposition 3.6. *Let M^{n+1} be an asymptotically harmonic CH-manifold. For any $u \in SM$ we have*

- i. *if M satisfies $K_M \leq -a^2$, then $\text{Ric}_M(u, u) \geq -n^2h^2 + n(n-1)a^2$;*
- ii. *if M satisfies $K_M \geq -b^2$, then $\text{Ric}_M(u, u) \leq -n^2h^2 + n(n-1)b^2$.*

As a consequence, we have the following characterization of constant curvature spaces :

Corollary 3.7. *Let M^{n+1} be an asymptotically harmonic CH-manifold.*

- i. *if M satisfies $K_M \leq -a^2$ then $h \geq a$, and $h = a$ if and only if $M = \mathbb{M}^{n+1}(-a^2)$;*
- ii. *if M satisfies $K_M \geq -b^2$ then $h \leq b$, and $h = b$ if and only if $M = \mathbb{M}^{n+1}(-b^2)$.*

Proof. The curvature upper bound $K_M \leq -a^2$ implies $h \geq a$. If $h = a$, then proposition 3.6 gives $\text{Ric}_M \geq -na^2$, and since the Ricci curvature is a sum of n sectional curvatures which are not greater than $-a^2$, this implies that all the sectional curvatures are equal to $-a^2$. The proof is the same when assuming a curvature lower bound. $\square$

3.3 Growth of horospheres

It is well known that, on CH-manifolds with *pinched* curvature, horospheres have polynomial volume growth, whose degree depend on the bounds on the curvature (cf. [14]). We will now see that, under the asymptotic harmonicity assumption, an upper bound $K_M \leq -a^2 < 0$ is enough to estimate from above the polynomial growth of horospheres.

Let H_ξ be a horosphere centered in some point at infinity ξ , let b_ξ be the Busemann function vanishing on H_ξ , and let g_0 be the Riemannian metric induced on H_ξ . For each $t \in \mathbb{R}$, there is a natural diffeomorphism $\varphi_t : H_\xi \rightarrow H_\xi(t)$ defined by $\varphi_t(x) = \exp_x(t\nabla b_\xi)$, which in turns induces a diffeomorphism

$$\Phi \begin{cases} \mathbb{R} \times H_\xi(0) & \rightarrow M \\ (t, x) & \mapsto \varphi_t(x) \end{cases}$$

In these “horospherical” coordinates (t, x) , the metric of M reads $g = dt^2 + g_t$, where $g_t = \varphi_t^* g_{H_\xi(t)}$ and $g_{H_\xi(t)}$ is the induced Riemannian metric of $H_\xi(t)$.

When assuming a sectional curvature upper bound $K \leq -a^2$, the map φ_t increases the distance for $t > 0$ and decreases the distance if $t < 0$. In fact, as a consequence of comparison theorem for Jacobi fields, we have that all the eigenvalues of $d\varphi_t$ are greater than or equal to e^{at} if $t > 0$, and less than or equal to e^{at} if $t < 0$ (cf. [12]).

Now, it is a standard fact that the mean curvature gives the derivative of the volume form of a submanifold under a deformation. In our setting, if $dv_t = J_t(x)dv_0$ is the volume form of the metric g_t and $J_t(x)$ is the density of dv_t

with respect to dv_0 , we have $J'_t = nh_t J_t$, where h_t is the mean curvature of $H_\xi(t)$. By asymptotic harmonicity, we deduce that $dv_t = e^{nht} dv_0$ for all t ; therefore, in horocyclic coordinates the volume form of M reads $dv_M = e^{nht} dt dv_0$.

On the other hand, by theorem 3.3, the volume entropy of M is nh : heuristically, this means that the exponential rate of the volume growth of M comes from the behaviour of the volume form in the $\mathbb{R}$ direction, and that the volume growth of the slices $H_\xi(t)$ should be subexponential. Namely:

Theorem 3.8. *Let M^{n+1} be an asymptotically harmonic CH-manifold with sectional curvature upper bound $K_M \leq -a^2 < 0$. Then, there exists a constant C (depending only on n , a and h) such that, for any horosphere H of M , the balls of H satisfy $\text{vol}(B_x^H(r)) \leq Cr^{\frac{nh}{a}}$ for all $r > 0$.*

Proof. Let $H = H_\xi$ be a horosphere centered in ξ . For any $u, v \in TH$, Gauss equation implies that $K_H(u, v) = K_M(u, v) + \mathcal{A}_\xi(u, u)\mathcal{A}_\xi(v, v) - \mathcal{A}_\xi(u, v)^2$, where K_H and K_M are the sectional curvatures of H and M respectively, and $\mathcal{A}_\xi$ is the second fundamental form of H . Taking the trace with respect to v we get

$$\begin{aligned} \text{Ric}_H(u, u) &= \text{Ric}_M(u, u) - K_M(u, u) + nh\mathcal{A}_\xi(u, u) - \sum_i \mathcal{A}_\xi(u, e_i)^2 \\ &\geq \text{Ric}_M(u, u) - |\mathcal{A}_\xi|^2 \geq -2n^2h^2 + 2n(n-1)a^2 \end{aligned}$$

where the last inequality comes from (3.2) and Proposition 3.6. Therefore, by Bishop's comparison theorem, there exists a constant C (depending only on n , a and h) such that, for any x in H we have $\text{Vol}(B_x^H(1)) \leq C$.

Let now $x \in H$ and consider the map $\varphi_{-t} : H \rightarrow H_\xi(-t)$ defined above, for $t > 0$. As $K_M \leq -a^2$, we have $\varphi_{-t}(B_x^H(r)) \subset B_{\varphi_{-t}(x)}^{H_\xi(-t)}(e^{-at}r)$. Moreover, as $dv_{-t} = e^{-nht} dv_0$, we have $\text{vol}(\varphi_{-t}(B_x^H(r))) = e^{-nht} \text{vol}(B_x^H(r))$; so, choosing $t = \frac{\ln r}{a}$ we obtain

$$\text{vol}(B_x^H(r)) \leq e^{nh\frac{\ln r}{a}} \text{vol}(B_{\varphi_{-t}(x)}^{H_\xi(-t)}(1)) \leq Cr^{\frac{nh}{a}}$$

□

Remark 3.9. This theorem proves that the degree of the polynomial volume growth of the horospheres is bounded above by $\frac{nh}{a}$. This upper bound is sharp, as it is the degree of the volume growth of the horospheres in the hyperbolic space (the horospheres being Euclidean in that case). Note that the upper bound is also sharp for the rank one symmetric spaces.

Remark 3.10. Using a similar proof, it is easy to see that the lower bound $-b^2 \leq K_M \leq 0$ gives a lower bound on the volume growth of the horospheres, namely $\text{vol}(B_x^H(r)) \geq Cr^{\frac{nb}{b}}$. The proof is left to the reader.

3.4 The mean value property

Harmonic manifolds are characterized by the fact that the harmonic functions have the mean value property : for any harmonic function F and any $R > 0$,

$$F(x_0) = \frac{1}{\text{vol}(S_{x_0}(R))} \int_{S_{x_0}(R)} F(x) dv_{S_{x_0}(R)}$$

This can be proved by taking the derivative of the right-hand side of the above equality, and by observing that it vanishes for any harmonic function F if and only if the spheres have constant mean curvature.

In the following theorem we prove that harmonic functions on an asymptotically harmonic manifold satisfy a mean value property, where, naturally, the mean is taken on horospheres. As the horospheres are non-compact, the mean on an horosphere is obtained as the limit of the means on an exhaustion. The computations of these horospherical means are very similar to those in [14].

Theorem 3.11. *Let M^{n+1} be an asymptotically harmonic manifold with sectional curvature upper bound $K_M \leq -a^2 < 0$, and let F be a function which is continuous on $M \cup \partial_\infty M$ and harmonic on M .*

For any $\xi \in \partial_\infty M$, any horosphere H_ξ centered in ξ , and any $x \in H_\xi$, there exists a sequence $(r_j)_{j \in \mathbb{N}}$ tending to $+\infty$ such that

$$\lim_{j \rightarrow \infty} \frac{1}{\text{Vol}(B_x^{H_\xi}(r_j))} \int_{B_x^{H_\xi}(r_j)} F dv_{H_\xi} = F(\xi)$$

where $B_x^{H_\xi}(R)$ denote the ball in H_ξ centered in x of radius R .

Proof. Let H_ξ be a horosphere centered in some point at infinity ξ , and let $\varphi_t : H_\xi \rightarrow H_\xi(t)$ be the diffeomorphism defined in §3.3.

Choose $x \in H_\xi$. Because H_ξ has polynomial volume growth, there exists a sequence $(r_j)_{j \in \mathbb{N}}$ tending to $+\infty$ such that

$$\lim_{j \rightarrow \infty} \frac{\text{vol}(\partial B_x^{H_\xi}(r_j))}{\text{Vol}(B_x^{H_\xi}(r_j))} = 0.$$

For $t \in \mathbb{R}$ and $j \in \mathbb{N}$, let $\Omega_{j,t} = \varphi_t(B_x^{H_\xi}(r_j))$. As pointed out in §3.3, we have $\text{Vol}(\Omega_{j,t}) = e^{nht} \text{Vol}(B_x^{H_\xi}(r_j))$. Moreover, the boundary of $\Omega_{j,t}$ satisfy

$$\frac{d}{dt} \text{vol}(\partial \Omega_{j,t}) = -(n-1) \int_{\partial \Omega_{j,t}} \langle \vec{k}_{j,t}, \frac{\partial}{\partial t} \rangle$$

where $\vec{k}_{j,t}$ is the mean curvature vector of $\partial \Omega_{j,t}$ (seen as a submanifold of M). Taking an orthonormal basis $(e_1, \dots, e_{n-1})$ of $T\partial \Omega_{j,t}$ and $\eta_{j,t}$ its exterior unit normal in $H_\xi(t)$ we have

$$-(n-1) \langle \vec{k}_{j,t}, \frac{\partial}{\partial t} \rangle = \sum_{i=1}^{n-1} \langle D_{e_i}^M \frac{\partial}{\partial t}, e_i \rangle = nh - \langle D_{\eta_{j,t}}^M \frac{\partial}{\partial t}, \eta_{j,t} \rangle \leq nh - a$$

where the last inequality comes from the curvature upper-bound on M . Therefore we have $\frac{d}{dt} \text{vol}(\partial \Omega_{j,t}) \leq (nh - a) \text{vol}(\partial \Omega_{j,t})$, and integrating this inequality we get $\text{vol}(\partial \Omega_{j,t}) \leq e^{(nh-a)t} \text{vol}(\partial \Omega_{j,0})$ and

$$\frac{\text{vol}(\partial \Omega_{j,t})}{\text{Vol}(\Omega_{j,t})} \leq e^{-at} \frac{\text{vol}(\partial B_x^{H_\xi}(r_j))}{\text{Vol}(B_x^{H_\xi}(r_j))}. \quad (3.4)$$

Consider now

$$g_j(t) = \frac{1}{\text{Vol}(\Omega_{j,t})} \int_{\Omega_{j,t}} F dv_t \quad (3.5)$$

where dv_t is the volume form of $H_\xi(t)$ and F a function which is continuous on $M \cup \partial_\infty M$ and harmonic on M . In particular, F is bounded. Using the fact that horospheres have constant mean curvature, we have

$$g'_j(t) = \frac{1}{\text{Vol}(\Omega_{j,t})} \int_{\Omega_{j,t}} \langle \nabla F, \frac{\partial}{\partial t} \rangle dv_t \quad (3.6)$$

and

$$g''_j(t) = \frac{1}{\text{Vol}(\Omega_{j,t})} \int_{\Omega_{j,t}} (\text{Hess}^M F)(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}) dv_t \quad (3.7)$$

Using proposition 1.1 and the fact that F is harmonic in M we get

$$(\text{Hess}^M F)(\frac{\partial}{\partial t}, \frac{\partial}{\partial t}) = -\text{tr}((\text{Hess}^{H_\xi(t)} F)|_{T_{H_\xi(t)}}) = \Delta^{H_\xi(t)} f + nh \langle \nabla F, \frac{\partial}{\partial t} \rangle$$

where f is the restriction of F to $H_\xi(t)$. Equation (3.7) gives

$$\begin{aligned} g''_j(t) - nhg'_j(t) &= \frac{1}{\text{Vol}(\Omega_{j,t})} \int_{\Omega_{j,t}} \Delta^{H_\xi(t)} f dv_t \\ &= -\frac{1}{\text{Vol}(\Omega_{j,t})} \int_{\partial\Omega_{j,t}} \langle \nabla F, \eta_{j,t} \rangle dv_t. \end{aligned} \quad (3.8)$$

As Ric_M is bounded from below and F is bounded on M , using Yau's gradient estimate for harmonic functions [25], there exists a constant C (depending on n, a, h and $\|F\|_\infty$) such that $|\nabla F| \leq C$ on M . Therefore, using (3.4), the right-hand side of (3.8) satisfies

$$\left| \frac{1}{\text{Vol}(\Omega_{j,t})} \int_{\partial\Omega_{j,t}} \langle \nabla F, \eta_{j,t} \rangle dv_t \right| \leq Ce^{-at} \frac{\text{vol}(\partial B_x^{H_\xi}(r_j))}{\text{Vol}(B_x^{H_\xi}(r_j))}$$

and tends uniformly to zero on bounded intervals when j tends to $+\infty$. In particular, it implies that, on bounded intervals, the C^0 norms of the functions g''_j are uniformly bounded. The fact that F is bounded and Yau's gradient estimate also imply that the C^0 norms of the functions g_j and g'_j are uniformly bounded, and, using Arzela-Ascoli convergence theorem, we have that, up to a subsequence, $(g_j)_{j \in \mathbb{N}}$ tends in C^1 topology to a function g .

Moreover, multiplying (3.8) by a test function, integrating by part and letting j tend to $+\infty$ we find that, in the sense of distributions, g is a solution of

$$g''(t) - nhg'(t) = 0.$$

Therefore, by classical regularity theory, g is smooth and $g'(t) = g'(0)e^{nht}$. Since g' is bounded on $\mathbb{R}$ we must have $g' \equiv 0$ and g is constant.

For any neighbourhood U of ξ (for the cone topology) there exist t such that the horosphere $H_\xi(t)$ is contained in U . By continuity of F on $M \cup \partial_\infty M$ and by the definition of g_j , the value $g_j(t)$ can be made arbitrary close to $F(\xi)$ (for any j). Therefore we have $g(t) = F(\xi)$ for any $t \in \mathbb{R}$, and $g(0) = F(\xi)$ gives the result. $\square$

Remark 3.12. It would be better to have a similar result without taking a sequence of radii tending to infinity, that is to have

$$\lim_{r \rightarrow \infty} \frac{1}{\text{Vol}(B_x^{H_\xi}(r))} \int_{B_x^{H_\xi}(r)} F dv_{H_\xi} = F(\xi).$$

For the proof to work in that case, one need to have $\lim_{r \rightarrow \infty} \frac{\text{vol}(\partial B_x^{H_\varepsilon}(r))}{\text{Vol}(B_x^{H_\varepsilon}(r))} =$

0. However, from the polynomial volume growth of horospheres one only get $\liminf_{r \rightarrow \infty} \frac{\text{vol}(\partial B_x^{H_\varepsilon}(r))}{\text{Vol}(B_x^{H_\varepsilon}(r))} = 0$.

4 Asymptotic behaviour of the volume form

In the previous section, in order to compute the entropy, we integrated the inequalities of lemma 2.6 on spheres. But since these inequalities hold pointwise, we can try to determine the asymptotic behaviour of the volume form at least in a fixed direction. Actually, let $\theta_x(u, r)$ be the density of the volume form of M in normal coordinates centered in some point x ; so the volume form reads $dv_M = \theta_x(u, r)dv_{S_x M}dr$, where $dv_{S_x M}$ is the volume form of $S_x M$.

Harmonic manifolds are characterized by the fact that $\theta_x(u, r)$ only depends on r . In this section we give a characterization of asymptotically harmonic manifolds in term of the asymptotic behaviour of $\theta_x(u, r)$:

Theorem 4.1. *Let M be a CH-manifold with $K_M \leq -a^2 < 0$ and entropy E . M is asymptotically harmonic if and only if there exists a positive function $\tau : SM \rightarrow \mathbb{R}_+$ such that $\theta_x(u, r)$ is uniformly equivalent to $\tau(u)e^{Er}$ for $r \rightarrow \infty$.*

“Uniformly equivalent” here means that the quotient of $\theta_x(u, r)$ by $\tau(u)e^{Er}$ converges to 1 for $r \rightarrow \infty$, uniformly with respect to $u \in SM$. This result will be consequence of the three propositions proved in the following subsections.

A Riemannian manifold is harmonic if and only if the density function only depends on r . As an asymptotic analogue, one would expect that $\lim_{r \rightarrow \infty} \frac{\theta(u, r)}{e^{Er}}$ does not depend on u , and thus that $\tau(u)$ be constant on SM . In proposition 4.3 we prove it holds under the restrictive assumption that DR_M is bounded.

4.1 The asymptotic volume-density function τ

The function θ_x is related to the mean curvature h_x of spheres centered in x of radius r by the formula

$$\frac{\theta'_x(u, r)}{\theta_x(u, r)} = nh_x(\exp_x(ru)) \quad (4.1)$$

where θ'_x denotes the derivative of θ_x with respect to r .

In what follows, we shall often write for short the point $\exp_x(ru)$ as (u, r) to avoid cumbersome notations; moreover, we will regard $\theta_x(u, r)$ as a function on $SM \times \mathbb{R}$, so we can drop the index x .

Using lemma 2.6 we get the following result :

Proposition 4.2. *Let M be a CH-manifold with curvature $K_M \leq -a^2 < 0$. If M is asymptotically harmonic, then there exists a bounded, positive function $\tau : SM \rightarrow \mathbb{R}_+$ such that*

$$\forall u \in SM \quad \left| \frac{\theta(u, r)}{\tau(u)e^{nr}} - 1 \right| \leq \varepsilon(r)$$

for an explicit function $\varepsilon(r)$ only depending on a and n such that $\lim_{r \rightarrow \infty} \varepsilon(r) = 0$.

Proof. As $\frac{\theta'(u,r)}{\theta(u,r)} = h_x(u,r)$, taking traces in (2.8) yields:

$$0 \leq \frac{\theta'(u,r)}{\theta(u,r)} - nh \leq na(\coth(ar) - 1),$$

that is

$$0 \leq \frac{d}{dr} [\ln(\theta(u,r)e^{-nhr})] \leq na [\coth(ar) - 1].$$

The first inequality implies that $\theta(u,r)e^{-nhr}$ is nondecreasing with respect to r . On the other hand, integrating the second one gives

$$\theta(u,r)e^{-nhr} \leq \theta(u,s)e^{-nhs} e^{na \int_s^r (\coth(at)-1)dt} \quad (4.2)$$

and, as $\int_s^\infty (\coth(at) - 1)dt$ is finite, we deduce that $\theta(u,r)e^{-nhr}$ is bounded, which implies that $\lim_{r \rightarrow \infty} \theta(u,r)e^{-nhr}$ exists. Therefore we can define the function τ on the unitary tangent bundle as

$$\tau(u) = \lim_{r \rightarrow \infty} \theta(u,r)e^{-nhr} \quad (4.3)$$

Moreover, as $\theta(u,r)e^{-nhr}$ is nondecreasing we have

$$\forall r > 0 \quad 0 < \theta(u,r)e^{-nhr} \leq \tau(u) \quad (4.4)$$

Again from (4.2), subtracting $\theta(u,s)e^{-nhs}$ and letting $r \rightarrow \infty$ we deduce

$$\tau(u) - \theta(u,s)e^{-nhs} \leq \theta(u,s)e^{-nhs} (e^{na \int_s^\infty (\coth(at)-1)dt} - 1)$$

As $\theta(u,s)e^{-nhs}$ is nondecreasing the left-hand side is nonnegative, so by (4.4),

$$\left| \frac{\theta(u,r)}{\tau(u)e^{nhr}} - 1 \right| \leq e^{na \int_r^\infty (\coth(at)-1)dt} - 1$$

The right-hand side is uniformly bounded from above and tends to 0 when r tends to infinity, which concludes the proof. $\square$

4.2 Properties of the function τ

First, we remark that the function τ is bounded : for any $u \in SM$ we have $\tau(u) \leq \frac{1}{(2a)^n}$. In fact, from equation (4.2) we obtain

$$\theta(u,r)e^{-nhr} \leq \frac{\theta(u,s)e^{-nhs}}{\sinh^n(as)} e^{-na(r-s)} \sinh^n(ar).$$

Letting s tend to 0, as $\frac{\theta(u,s)}{s^n} \rightarrow 1$, we deduce

$$\theta(u,r)e^{-nhr} \leq \frac{e^{-nar} \sinh^n(ar)}{a^n},$$

and for $r \rightarrow \infty$ we get $\tau(u) \leq \frac{1}{(2a)^n}$.

Therefore, proposition 4.2 implies

$$|\theta(u,r)e^{-nhr} - \tau(u)| \leq \frac{1}{(2a)^n} \varepsilon(r)$$

and the function τ is the uniform limit of $\theta(u, r)e^{-nhr}$; as the convergence is uniform, the function τ is continuous on SM . Moreover, *as soon as $\theta(u, r)e^{-nhr}$ has a limit*, this limit can be expressed in terms of Jacobi tensors. This was used to study the asymptotic behaviour of the volume on harmonic manifolds (cf. [16, 6, 11]). Using this approach we get more information on the function τ .

Proposition 4.3. *Let M be an asymptotically harmonic CH-manifold with curvature $K_M \leq -a^2 < 0$. Then:*

i. $\tau : SM \rightarrow \mathbb{R}_+$ is invariant by the geodesic flow and flip invariant, i.e. :

- $\tau(\dot{\gamma}(t))$ is constant for any geodesic γ ;
- $\tau(v) = \tau(-v)$ for all $v \in SM$.

ii. if DR_M is bounded on M , then τ is constant on SM ;

iii. $\tau \geq \frac{1}{(2h)^n}$, with equality if and only if the curvature is constant.

Proof. A Jacobi tensor along a geodesic γ is a smooth family $J(t)$ of endomorphisms of $\dot{\gamma}(t)^\perp$ satisfying the Jacobi equation $J''(t) + R(t)J(t) = 0$, where $R(t)$ is defined from the Riemann tensor by $R(t)u = R(\dot{\gamma}(t), u)\dot{\gamma}(t)$. Then, applying J to any parallel vector field $V(t)$ along γ gives a Jacobi vector field $J(t)V(t)$.

Let $v \in S_x M$ and $\gamma(t) = \exp_x(tv)$, and consider the Jacobi tensor J_v along γ defined by $J_v(0) = 0$ and $J'_v(0) = \text{Id}$. It is well known that $J'_v(r)J_v^{-1}(r)$ gives the shape operator $A_x(v, r)$ of the sphere $S_x(r)$ at $\exp_x(rv)$ (with respect to the inner normal to the sphere), and that $\theta(v, r) = \det(J_v(r))$.

For $r > 0$, let $U_{v,r}, S_{v,r}$ be the Jacobi tensors on γ defined by $U_{v,r}(-r) = 0$, $U_{v,r}(0) = S_{v,r}(0) = \text{Id}$ and $S_{v,r}(r) = 0$. The unstable and stable Jacobi tensors at v are defined by $U_v = \lim_{r \rightarrow \infty} U_{v,r}$ and $S_v = \lim_{r \rightarrow \infty} S_{v,r}$. As $U'_{v,r}(0) = J'_{\dot{\gamma}(-r)}(r)J_{\dot{\gamma}(-r)}^{-1}(r)$ is the shape operator of the sphere $S_{\dot{\gamma}(-r)}(r)$ at x , it follows that $U'_v(0)$ is shape operator at x of the horosphere centered in $\xi_- = \lim_{r \rightarrow \infty} \gamma(-r)$. In a similar way, we have that $-S'_v(0)$ is the shape operator at x of the horosphere centered in $\xi_+ = \lim_{r \rightarrow \infty} \gamma(r)$. Since M is asymptotically harmonic, we have $\text{tr}(U'_v(0)) = nh$; following the proof of Corollary 2.5 of [16] we get

$$\theta(v, t)e^{-nht} = \frac{1}{\det(U'_v(0) - S'_{v,t}(0))},$$

which, taking the limit for $t \rightarrow \infty$, gives

$$\tau(v) = \frac{1}{\det(U'_v(0) - S'_v(0))}. \quad (4.5)$$

The proposition then follows from this expression of $\tau(v)$.

First, as $U'_v(0)$ and $-S'_v(0)$ are the shape operators of the horospheres centered in ξ_- and ξ_+ , relative to their respective inner normals, it is clear that τ is flip invariant. The invariance by the geodesic flow is just lemma 2.2 in [11].

To prove the second point, let us first show that $\tau(u) = \tau(v)$ when $u, v \in SM$ point towards the same boundary point $\xi \in \partial_\infty M$, i.e. $\lim_{s \rightarrow +\infty} \gamma_u(s) = \lim_{s \rightarrow +\infty} \gamma_v(s)$. By the invariance of τ under the geodesic flow, we may as well

assume that u and v are normal to the same horosphere, so $d(\gamma_u(t), \gamma_v(t)) \leq c_1 e^{-at}$ for all $t > 0$. For any $r, t > 0$ we have

$$|\tau(u) - \tau(v)| \leq |\tau(u) - \theta(\dot{\gamma}_u(t), r)e^{-nhr}| + |\theta(\dot{\gamma}_u(t), r) - \theta(\dot{\gamma}_v(t), r)|e^{-nhr} \\ + |\tau(v) - \theta(\dot{\gamma}_v(t), r)e^{-nhr}|,$$

and using the invariance of τ by the geodesic flow and Proposition 4.2 we get

$$|\tau(u) - \tau(v)| \leq (\tau(u) + \tau(v))\varepsilon(r) + |\theta(\dot{\gamma}_u(t), r) - \theta(\dot{\gamma}_v(t), r)|e^{-nhr}. \quad (4.6)$$

For $s \in]0, r]$, let $h_{u,t}(s)$ (resp. $h_{v,t}(s)$) be the mean curvature, at the point $\gamma_u(t+s)$ (resp. at $\gamma_v(t+s)$), of the sphere of radius s centered in $\gamma_u(t)$ (resp. $\gamma_v(t)$). Following the Lemma 2.3 in [11], we will use comparison theory for Riccati equation to estimate $|h_{u,t}(s) - h_{v,t}(s)|$. We choose orthonormal parallel basis $e_{u,i}(s)$ of $\dot{\gamma}_u(t+s)^\perp$ and $e_{v,i}(s)$ of $\dot{\gamma}_v(t+s)^\perp$ such that, for any i , $d(e_{u,i}(s), e_{v,i}(s)) \leq c_2 e^{-a(t+s)}$ in SM , for some constant c_2 . Let $A_{u,t}(s)$ and $A_{v,t}(s)$ be the matrices of the second fundamental forms of the spheres of radius s centered in $\gamma_u(t), \gamma_v(t)$ in these basis. They satisfy the Riccati equations $A'_{u,t}(s) + A_{u,t}^2(s) + R_{u,t}(s) = 0$ and $A'_{v,t}(s) + A_{v,t}^2(s) + R_{v,t}(s) = 0$, where $R_{u,t}(s)$ is the matrix of the endomorphism $R(\dot{\gamma}_u(t+s), \cdot)\dot{\gamma}_u(t+s)$, and analogously for $R_{v,t}(s)$. Because of the assumption on DR_M , we have that the tensor $r(s) = R_{u,t}(s) - R_{v,t}(s)$ satisfies

$$|r(s)| \leq C_3 e^{-a(t+s)}. \quad (4.7)$$

Consider now $B(s) = A_{u,t}(s) - A_{v,t}(s)$ and $Q(s) = \frac{1}{2}(A_{u,t}(s) + A_{v,t}(s))$. From the Riccati equations we have that B is solution of

$$B'(s) + B(s)Q(s) + Q(s)B(s) + r(s) = 0.$$

A direct computation shows that for any $0 < \varepsilon < s$ we have the formula

$$B(s) = {}^t C(s) \left[{}^t C(\varepsilon)^{-1} B(\varepsilon) C(\varepsilon)^{-1} - \int_\varepsilon^s {}^t C(\zeta)^{-1} r(\zeta) C(\zeta)^{-1} d\zeta \right] C(s) \quad (4.8)$$

where $C(s)$ is a solution of $C'(s) = -C(s)Q(s)$. In particular, because of the curvature upper bound we have $Q(s) \geq a\text{Id}$ hence, for any $0 < \varepsilon < s$, $|C(\varepsilon)^{-1} C(s)| \leq e^{-a(s-\varepsilon)}$. Plugging this estimate and (4.7) in the formula (4.8) we get

$$|B(s)| \leq |B(\varepsilon)|e^{-2a(s-\varepsilon)} + c_4 e^{-a(t+s)}$$

Since both $A_{u,t}(s)$ and $A_{v,t}(s)$ behave, for $s \rightarrow 0$, as $\frac{1}{s}\text{Id} + o(1)$ we have $\lim_{\varepsilon \rightarrow 0} B(\varepsilon) = 0$; therefore we deduce that $|B(s)| \leq c_4 e^{-a(t+s)}$ and, taking the trace,

$$|h_{u,t}(s) - h_{v,t}(s)| \leq c_5 e^{-a(t+s)}$$

for some constant c_5 . By the expression (4.1) for h_x , integrating on $[0, r]$ yields

$$-c_6(1 - e^{-ar})e^{-at} \leq \ln \frac{\theta(\dot{\gamma}_u(t), r)}{\theta(\dot{\gamma}_v(t), r)} \leq c_6(1 - e^{-ar})e^{-at}$$

With these inequalities we can bound the last term of (4.6):

$$\begin{aligned} & \theta(\dot{\gamma}_v(t), r) e^{-nhr} [\exp(-c_6(1 - e^{-ar})e^{-at}) - 1] \\ & \leq |\theta(\dot{\gamma}_u(t), r) - \theta(\dot{\gamma}_v(t), r)| e^{-nhr} \\ & \leq \theta(\dot{\gamma}_v(t), r) e^{-nhr} [\exp(c_6(1 - e^{-ar})e^{-at}) - 1] \quad (4.9) \end{aligned}$$

Choosing r large enough, the term $\varepsilon(r)$ in (4.6) can be made arbitrary small; for this value of r , $\theta(\dot{\gamma}_v(t), r) e^{-nhr}$ stays close to $\tau(v)$ for all t by proposition 4.2, and the above estimate (4.9) implies that we can choose t large enough to make also the last term of (4.6) arbitrary small. Therefore $\tau(u) = \tau(v)$.

Consider now any vector $u, v \in SM$, and let σ be a geodesic such that $\lim_{s \rightarrow -\infty} \sigma(s) = \lim_{s \rightarrow +\infty} \gamma_u(s)$ and $\lim_{s \rightarrow +\infty} \sigma(s) = \lim_{s \rightarrow +\infty} \gamma_v(s)$. From the above computations we must have $\tau(v) = \tau(\dot{\sigma}(0))$ and $\tau(u) = \tau(-\dot{\sigma}(0))$, so by flip invariance we get $\tau(u) = \tau(v)$. Therefore τ is constant.

The third point is similar to Corollary 2.6 in [16]. As $U'_v(0) - S'_v(0)$ is a positive symmetric matrix, the arithmetic-geometric inequality gives

$$\det(U'_v(0) - S'_v(0))^{\frac{1}{n}} \leq \frac{1}{n} \text{tr}(U'_v(0) - S'_v(0)) = 2h$$

and the inequality follows. The case of equality follows, as in the proof of Corollary 2.6 in [16], from the fact that $s \mapsto U'_{\dot{\gamma}(s)}(0)$ and $s \mapsto S'_{\dot{\gamma}(s)}(0)$ satisfy the Riccati equation, and because $U'_{\dot{\gamma}(s)}(0) - S'_{\dot{\gamma}(s)}(0) = \frac{2h}{n-1} Id$. $\square$

Remark 4.4. The second point is very close to Lemma 2.3 and Corollary 2.1 of [11], but, in our proof, we don't need any lower bound on the curvature and rather use Proposition 4.2.

4.3 Characterization of asymptotic harmonicity

Proposition 4.2 says that the volume form of M has purely exponential growth, with isotropic exponential rate (and is asymptotically perfectly isotropic when DR_M is bounded). In fact this is a characterization of asymptotic harmonicity:

Proposition 4.5. *Let M be a CH-manifold with $K_M \leq -a^2 < 0$ and entropy E . If there exists a positive function $\tau : SM \rightarrow \mathbb{R}$ such that $\theta(u, r)$ is uniformly equivalent to $\tau(u)e^{Er}$ for $r \rightarrow \infty$, then M is asymptotically harmonic.*

Remark 4.6. Notice that, together with Benoist-Foulon-Labourie and Besson-Courtois-Gallot characterization of cocompact asymptotically harmonic spaces, Proposition 4.5 shows that if a CH-manifold with compact quotients has volume form which is (uniformly) equivalent to a function $\tau(u)e^{ER}$, then it is a ROSS.

Proof. Let $\gamma(t)$ be a geodesic of M with $\lim_{t \rightarrow -\infty} \gamma(t) = \xi \in \partial_\infty M$, and let $h(t)$ be the mean curvature at $\gamma(t)$ of the horosphere $H_\xi(t)$ centered in ξ and passing through $\gamma(t)$. We shall prove that the function $h(t)$ is constant.

Let $r < R$ be two real numbers, and choose $s > -r$. For any $t \in [r, R]$, we use Lemma 2.6 to compare the second fundamental forms of $H_\xi(t)$ and $S_{\gamma(-s)}(t+s)$ at $\gamma(t)$. Taking the trace in (2.8), we have

$$0 \leq \frac{\theta'(\dot{\gamma}(-s), t+s)}{\theta(\dot{\gamma}(-s), t+s)} - nh(t) \leq na(\coth(a(t+s)) - 1)$$

and integrating on $[r, R]$ with respect to t we get

$$0 \leq \ln \frac{\theta(\dot{\gamma}(-s), R+s)}{\theta(\dot{\gamma}(-s), r+s)} - n \int_r^R h(t) dt \leq \ln \left(\frac{\sinh^n(a(R+s))}{\sinh^n(a(r+s))} e^{-na(R-r)} \right) \quad (4.10)$$

The right-hand side tends to 0 when s tends to infinity. Moreover, by hypothesis we have $|\frac{\theta(\dot{\gamma}(-s), R+s)}{\tau(\dot{\gamma}(-s))e^{E(R+s)}} - 1| \leq \varepsilon(R+s)$ with $\lim_{s \rightarrow \infty} \varepsilon(R+s) = 0$, and we get

$$\lim_{s \rightarrow \infty} \frac{\theta(\dot{\gamma}(-s), R+s)}{\tau(\dot{\gamma}(-s))e^{Es}} = e^{ER}.$$

Analogously, we find $\lim_{s \rightarrow \infty} \frac{\theta(\dot{\gamma}(-s), r+s)}{\tau(\dot{\gamma}(-s))e^{Es}} = e^{Er}$, so letting s tend to infinity in (4.10) we obtain

$$E(R-r) - n \int_r^R h(t) dt = 0.$$

Therefore $\int_r^R (E - nh(t)) dt = 0$ for all $r < R$, from which we deduce that $h(t) = \frac{E}{n}$ for all $t \in \mathbb{R}$, and M is asymptotically harmonic. $\square$

5 Margulis function and measures at infinity

In this last section, we assume that M is a asymptotically harmonic CH-manifold with pinched curvature $-b^2 \leq K_M \leq -a^2 < 0$, and h is always the mean curvature of the horospheres.

5.1 Visual and harmonic measures

There are two families of measures naturally defined on the ideal boundary of Cartan-Hadamard manifolds: the visual and harmonic measures.

To define the visual measures, consider the homeomorphism given by the “projection on $\partial_\infty M$ from x ”:

$$\phi_x : \begin{cases} S_x M & \rightarrow \partial_\infty M \\ u & \mapsto \phi_x(u) = \lim_{t \rightarrow \infty} \exp_x(tu) \end{cases}$$

The measure λ_x is the push-forward on $\partial_\infty M$ of the (normalized) Riemannian measure of $S_x M$.

On the other hand, the family of harmonic measures comes from the uniqueness of the solution to the Dirichlet problem at infinity (cf. [1]): given a continuous function f on $\partial_\infty M$, there exists a unique bounded harmonic function F on M such that $\lim_{x \rightarrow \xi} F(x) = \xi$. Then, it is a consequence of Riesz representation theorem that there exists a unique family of measures μ_x , $x \in M$, such that $F(x) = \int_{\partial_\infty M} f(\xi) d\mu_x(\xi)$.

Proposition 5.1. *Let M be an asymptotically harmonic CH-manifold with pinched curvature $-b^2 \leq K_M \leq -a^2 < 0$. For any $x, y \in M$ we have*

$$\frac{d\lambda_x}{d\lambda_y}(\xi) = \frac{\tau(\phi_y^{-1}(\xi))}{\tau(\phi_x^{-1}(\xi))} e^{-nh(b_\xi(x) - b_\xi(y))} \quad \text{and} \quad \frac{d\mu_x}{d\mu_y}(\xi) = e^{-nh(b_\xi(x) - b_\xi(y))}.$$

Proof. Consider the distance functions r_x and r_y to $x, y \in M$ respectively, and the sphere $S_x(t)$ centered in x of radius t . For t great enough, each geodesic ray from y intersect $S_x(t)$ at a unique point; for $v \in S_y M$, let $F_t(v)$ be the intersection point of the geodesic $s \mapsto \exp_y(sv)$ and $S_x(t)$.

The map $F_t : S_y M \rightarrow S_x(t)$ so defined is a diffeomorphism whose Jacobian is

$$\text{Jac}_v F_t = \frac{\theta(v, r_y(F_t(v)))}{\langle \nabla r_y(F_t(v)), \nabla r_x(F_t(v)) \rangle} \quad (5.1)$$

Now, let $U \subset \partial_\infty M$ be a measurable set with negligible boundary, and let $U_t = \{\exp_x(tu) \mid u \in \phi_x^{-1}(U)\}$ be the projection of U on $S_x(t)$ from x .

By definition of λ_x we have

$$\lambda_x(U) = \int_{\phi_x^{-1}(U)} d\sigma_x = \frac{1}{\text{vol}(S^n)} \int_{U_t} \frac{1}{\theta(P_t^{-1}(z), t)} dv_{S_x(t)}(z)$$

where $P_t(u) = \exp_x(tu)$ for $u \in S_x M$, $d\sigma_x$ is the normalized measure of $S_x M$, and $dv_{S_x(t)}$ the volume forms of $S_x(t)$.

By (5.1), we get

$$\lambda_x(U) = \int_{F_t^{-1}(U_t)} \frac{\theta(v, r_y(F_t(v)))}{\theta(P_t^{-1} \circ F_t(v), t)} \langle \nabla r_y(F_t(v)), \nabla r_x(F_t(v)) \rangle^{-1} d\sigma_y(v) \quad (5.2)$$

where $d\sigma_y$ is the normalized measure on $S_y M$.

Now we observe that, letting t tend to infinity, we have

- $\lim_{t \rightarrow \infty} P_t^{-1} \circ F_t(v) = \phi_x^{-1} \circ \phi_y(v)$;
- $\lim_{t \rightarrow \infty} \chi_{F_t^{-1}(U_t)} = \chi_{\phi_y^{-1}(U)}$ almost everywhere;
- $\lim_{t \rightarrow \infty} \langle \nabla r_y(F_t(v)), \nabla r_x(F_t(v)) \rangle = 1$

Moreover, from Theorem 4.2 we know that

$$\begin{aligned} \frac{\tau(v) - \varepsilon(r_y(F_t(v)))}{\tau(P_t^{-1} \circ F_t(v)) + \varepsilon(t)} e^{nh(r_y(F_t(v)) - t)} &\leq \frac{\theta_y(v, r_y(F_t(v)))}{\theta(P_t^{-1} \circ F_t(v), t)} \\ &\leq \frac{\tau(v) + \varepsilon(r_y(F_t(v)))}{\tau(P_t^{-1} \circ F_t(v)) - \varepsilon(t)} e^{nh(r_y(F_t(v)) - t)} \end{aligned}$$

By definition of Busemann function we have that $r_y(F_t(v)) - t$ converges, uniformly on $S_y M$, to $b_{\phi_y(v)}(y) - b_{\phi_y(v)}(x)$; so, as τ is continuous and bounded, by dominating convergence (5.2) yields

$$\begin{aligned} \lambda_x(U) &= \int_{\phi_y^{-1}(U)} \frac{\tau(v)}{\tau(\phi_x^{-1} \circ \phi_y(v))} e^{-nh(b_{\phi_y(v)}(x) - b_{\phi_y(v)}(y))} dv_{S_y M}(v) \\ &= \int_U \frac{\tau(\phi_y^{-1}(\xi))}{\tau(\phi_x^{-1}(\xi))} e^{-nh(b_\xi(x) - b_\xi(y))} d\lambda_y(\xi) \end{aligned}$$

which proves the first equality of the proposition.

The second equality follows from [1]: the relative densities of harmonic measures are given by the Poisson kernel, and, as $\Delta b_\xi = -nh$, by unicity of the Poisson kernel we have $\frac{d\mu_x}{d\mu_y}(\xi) = e^{-nh(b_\xi(x) - b_\xi(y))}$. $\square$

As a consequence of Theorem 4.3, we have that, *when the derivative of the Riemann tensor is bounded, the visual and harmonic measures class have the same relative densities.*

5.2 The Margulis function

For cocompact CH-manifolds Margulis introduced the function

$$m(x) = \lim_{r \rightarrow \infty} \text{vol}(S_x(r))e^{-Er}.$$

where E is the volume entropy of M . The main conjecture concerning this function is that it is constant if and only if M is a symmetric space, cf.[27, 15] for some related results. Theorem 4.2 allows us to define the Margulis function for asymptotically harmonic manifolds (even noncocompact) :

Proposition 5.2. *Let M be an asymptotically harmonic CH-manifold with $-b^2 \leq K \leq -a^2 < 0$. There exists a function $m : M \rightarrow \mathbb{R}_+$ such that*

$$\lim_{r \rightarrow \infty} \text{vol}(S_x(r))e^{-nhr} = m(x) \quad \text{and} \quad \lim_{r \rightarrow \infty} \text{vol}(B_x(r))e^{-nhr} = \frac{m(x)}{nh}$$

for any $x \in M$. Moreover, the function m is harmonic.

Proof. Let $V_x(r) = \text{Vol}(B_x(r))$ and $v_x(r) = \text{vol}(S_x(r))$, so $V'_x(r) = v_x(r)$. Since $v_x(r) = \int_{S_x M} \theta(u, r) du$, integrating (4.3) on $S_x M$, by monotone convergence we get the first equality with $m(x) = \int_{S_x M} \tau(u) du$.

Then, by Proposition 3.2, we have $V'_x(r) - nhV_x(r) \geq 0$, so $V_x(r)e^{-nhr}$ is increasing. As $V_x(r) = \int_0^r \int_{S_x M} \theta(u, s) duds$, Theorem 4.2 implies, for any $r \geq 1$,

$$V_x(r) \leq V_x(1) + \frac{m(x)}{nh}(e^{nhr} - e^{nh}) + \text{vol}(S^n) \int_1^r \varepsilon(s)e^{nhs} ds$$

from which we deduce that $V_x(r)e^{-nhr}$ is bounded; hence, it converges to some limit $l(x)$. As $V_x(r)e^{-nhr}$ is increasing and converging, there exists a sequence $r_k \rightarrow \infty$ such that

$$\begin{aligned} 0 &= \lim_{k \rightarrow \infty} \frac{d}{dr} \Big|_{r=r_k} (V_x(r)e^{-nhr}) \\ &= \lim_{k \rightarrow \infty} (v_x(r_k)e^{-nhr_k} - nhV_x(r_k)e^{-nhr_k}) \\ &= m(x) - nhl(x). \end{aligned}$$

Finally, to show that the Margulis function is harmonic, we write it using the visual measures :

$$m(x) = \int_{S_x M} \tau(u) du = \int_{\partial_\infty M} \tau(\phi_x^{-1}(\xi)) d\lambda_x(\xi)$$

Choosing a fixed point $x_0 \in M$, we get

$$m(x) = \int_{\partial_\infty M} \tau(\phi_{x_0}^{-1}(\xi)) e^{-nh(b_\xi(x) - b_\xi(x_0))} d\lambda_{x_0}(\xi)$$

and we are done, because $e^{-nh(b_\xi(x))}$ is harmonic. $\square$

References

- [1] M.T. Anderson and R. Schoen. Positive harmonic functions on complete manifolds of negative curvature. *Ann. of Math. (2)*, 121(3):429–461, 1985.
- [2] Y. Benoist, P. Foulon, and F. Labourie. Flots d’Anosov à distributions stable et instable différentiables. *J. Amer. Math. Soc.*, 5(1):33–74, 1992.
- [3] A.L. Besse. *Manifolds all of whose geodesics are closed*, volume 93 of *Ergebnisse der Mathematik und ihrer Grenzgebiete [Results in Mathematics and Related Areas]*. Springer-Verlag, Berlin, 1978. With appendices by D. B. A. Epstein, J.-P. Bourguignon, L. Bérard-Bergery, M. Berger and J. L. Kazdan.
- [4] G. Besson, G. Courtois, and S. Gallot. Entropies et rigidités des espaces localement symétriques de courbure strictement négative. *Geom. Funct. Anal.*, 5(5):731–799, 1995.
- [5] M. R. Bridson and A. Haefliger. *Metric spaces of non-positive curvature*, volume 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
- [6] C. Connell. Asymptotic harmonicity of negatively curved homogeneous spaces and their measures at infinity. *Comm. Anal. Geom.*, 8(3):575–633, 2000.
- [7] E. Damek and F. Ricci. A class of nonsymmetric harmonic Riemannian spaces. *Bull. Amer. Math. Soc. (N.S.)*, 27(1):139–142, 1992.
- [8] H. Donnelly. On the essential spectrum of a complete Riemannian manifold. *Topology*, 20(1):1–14, 1981.
- [9] P. Foulon and F. Labourie. Sur les variétés compactes asymptotiquement harmoniques. *Invent. Math.*, 109(1):97–111, 1992.
- [10] J. Heber. On harmonic and asymptotically harmonic homogeneous spaces. *Geom. Funct. Anal.*, 16(4):869–890, 2006.
- [11] J. Heber, G. Knieper, and H. Shah. Asymptotically harmonic spaces in dimension 3. *Proc. Amer. Math. Soc.*, 135(3):845–849 (electronic), 2007.
- [12] E. Heintze and H.C. Im Hof. Geometry of horospheres. *J. Differential Geom.*, 12:481–491, 1977.
- [13] H. Karcher. Riemannian comparison constructions. In *Global differential geometry*, volume 27 of *MAA Stud. Math.*, pages 170–222. Math. Assoc. America, Washington, DC, 1989.
- [14] L. Karp and N. Peyerimhoff. Horospherical means and uniform distribution of curves of constant geodesic curvature. *Math. Z.*, 231(4):655–677, 1999.
- [15] G. Knieper. Spherical means on compact Riemannian manifolds of negative curvature. *Differential Geom. Appl.*, 4(4):361–390, 1994.
- [16] G. Knieper. New results on noncompact harmonic manifolds. To appear in *Comment. Math. Helv.*, 2009.
- [17] H. Kumura. On the essential spectrum of the Laplacian on complete manifolds. *J. Math. Soc. Japan*, 49(1):1–14, 1997.
- [18] F. Ledrappier. Harmonic measures and Bowen-Margulis measures. *Israel J. Math.*, 71(3):275–287, 1990.
- [19] F. Ledrappier. Ergodic properties of the stable foliations. In *Ergodic Theory and Related Topics III*, volume 1514 of *Lecture Notes in Mathematics*, pages 131–145. Springer, 1992.
- [20] A. Ranjan and H. Shah. Harmonic manifolds with minimal horospheres. *J. Geom. Anal.*, 12(4):683–694, 2002.

- [21] A. Ranjan and H. Shah. Busemann functions in harmonic manifolds. *Geom. Dedicata*, 101:167–183, 2003.
- [22] M. Reed and B. Simon. *Methods of modern mathematical physics. I. Functional Analysis*. Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, second edition, 1980. Functional analysis.
- [23] V. Schroeder and H. Shah. On 3-dimensional asymptotically harmonic manifolds. *Arch. Math. (Basel)*, 90(3):275–278, 2008.
- [24] Z. I. Szabó. The Lichnerowicz conjecture on harmonic manifolds. *J. Differential Geom.*, 31(1):1–28, 1990.
- [25] S.T. Yau. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.*, 28:201–228, 1975.
- [26] C. Yue. Rigidity and dynamics around manifolds of negative curvature. *Math. Res. Lett.*, 2(1):123–147, 1994.
- [27] C. Yue. Brownian motion on Anosov foliations and manifolds of negative curvature. *J. Differential Geom.*, 41(1):159–183, 1995.

Philippe Castillon
 I3M (U.M.R. C.N.R.S. 5149)
 Dépt. des Sciences Mathématiques, CC 51
 Univ. Montpellier II
 34095 MONTPELLIER Cedex 5, France
 philippe.castillon@univ-montp2.fr

Andrea Sambusetti
 Istituto di Matematica G. Castelnuovo
 Università "La Sapienza" di Roma
 P.le Aldo Moro 5
 00185 Roma (Italy)
 sambuset@mat.uniroma1.it